



# Analysis of the correlation between sperm DNA integrity and conventional semen parameters in infertile men

## *İnfertil erkeklerde sperm DNA bütünlüğü ve konvansiyonel semen parametreleri arasındaki korelasyonun analizi*

Oya Sena Aydos<sup>1</sup>, Yunus Yükselten<sup>1</sup>, Fuat Kaplan<sup>1</sup>, Asuman Sunguroğlu<sup>1</sup>, Kaan Aydos<sup>2</sup>

### ABSTRACT

**Objective:** A male factor is responsible in approximately 30-40% of couples receiving infertility treatment. Routinely, such couples undergo semen analysis including parameters such as sperm count, motility and morphology. Generally, the analysis of sperm DNA damage, shown to have a significant clinical importance by many studies, is recognized as an advanced test that is not included in routine infertility tests. Intracytoplasmic sperm injection (ICSI) method, commonly employed in the current infertility treatment protocols, lowers the fertilization rate, however, fertilization can occur even with a damaged DNA which is known to pose a risk in the subsequent pregnancy period. The relation between sperm morphology and the degree of sperm DNA damage has not yet been understood clearly. In this study, we aimed to investigate the association between routine semen analysis and sperm DNA integrity assay, another advanced but costly method.

**Material and methods:** The degree of DNA damage was compared with the results of semen analysis, based on the WHO criteria, in 399 male patients who received comet assay for sperm DNA integrity. The statistical correlation analyses were performed with Windows SPSS statistical package program.

**Results:** Accordingly, the sperm DNA damage was found to be correlated with all 3 parameters (sperm count, forward motility, and morphology) examined by the semen analysis ( $p<0.001$ ). Total sperm DNA Damage Count (TDC) was 226, 216, and 210 arbitrary units (AU) in patients with a sperm count  $<15$  mil/mL, forward moving motility  $<32\%$ , and normal morphology  $<4\%$ , respectively. The difference with the normal individuals was statistically significant ( $p<0.001$ ).

**Conclusion:** In light of the comet assay results, higher degree of sperm DNA damage is associated with significant impairment of all seminal parameters.

**Keywords:** Male infertility; morphology; motility; sperm DNA damage.

### ÖZ

**Amaç:** İnfertilite tedavisi gören çiftlerin yaklaşık %30-40'ında erkek faktörü sorumludur. Bunların değerlendirilmesinde rutin olarak sperm konsantrasyonu, motilitesi ve morfolojisine bakılır. Klinik önemi çok sayıda çalışmada gösterilmiş olan sperm DNA hasar incelenmesi ise genelde ileri bir test olarak ele alınmakta ve rutin infertilite tetkikleri arasında yer almamaktadır. Oysa güncel infertilite tedavisinde çok yaygın olarak kullanılan, yüksek maliyeti yanı sıra birçok doğal bariyerin atlanmasına olanak tanıyan intrasitoplazmik sperm enjeksiyonu (ICSI) yönteminde, fertilizasyon oranı düşse de hasarlı DNA ile bile fertilizasyonun gerçekleşebildiği, bunun da gebeliğin devamında risk oluşturduğu bilinmektedir. Ancak ICSI sırasında sperm seçiminde sadece sperm morfoloji ve motilitesi dikkate alınmakta olup, DNA hasarına göre bir seçim teknik olarak yapılamamaktadır. Sperm morfolojisinin DNA içeriğindeki hasar oranları ile ilişkisi ise kesin olarak aydınlatılmış değildir. Biz bu çalışmada rutin semen analizinin daha ileri ancak maliyeti de yüksek bir tetkik olan sperm DNA bütünlüğü analiz sonuçlarıyla ilişkisini incelemeyi amaçladık.

**Gereç ve yöntemler:** Comet analizi ile sperm DNA bütünlüğü araştırılmış 399 erkek hastanın DNA hasar oranları, WHO kriterlerine göre değerlendirilmiş semen analiz sonuçları ile karşılaştırıldı ve Windows SPSS paket programı kullanılarak istatistiksel korelasyon analizleri yapıldı.

**Bulgular:** Buna göre sperm DNA hasarlarının semen analizinde bakılan her üç parametreyle (sayı, ileri motilite, morfoloji) anlamlı şekilde ilişkili olduğu saptandı ( $p<0.001$ ).

**Sonuç:** COMET assay ile tespit edilebildiği kadarıyla sperm DNA hasar oranlarındaki artış seminal parametrelerin hepsinde de anlamlı biçimde bozulma ile birlikte ortaya çıkmaktadır.

**Anahtar kelimeler:** Erkek infertilitesi; morfoloji; motilite; sperm DNA hasarı.

<sup>1</sup>Department of Medical Biology, Ankara University Faculty of Medicine, Ankara, Turkey

<sup>2</sup>Department of Urology, Ankara University Faculty of Medicine, Ankara, Turkey

**Submitted:**  
17.04.2015

**Accepted:**  
25.05.2015

**Correspondence:**  
Oya Sena Aydos  
E-mail: ksaydos@gmail.com

©Copyright 2015 by Turkish Association of Urology

Available online at  
www.turkishjournalofurology.com

## Introduction

In almost half of the infertile couples, a male factor is involved. These reproductive problems are routinely evaluated by conventional semen analysis. However, 15% of cases of male factor infertility have normal semen analysis which complicates a definitive diagnosis. There is a strong association between sperm genetic damage and infertility. The involvement of sperm DNA damage in infertility has been shown by many *in vitro* and *in vivo* studies. It has been found that spermatozoa with denatured and fragmented DNA have a significant relationship with impaired seminal parameters, being significantly higher in infertile males (25% and 27%, respectively) than in fertile ones (10% and 13%, respectively) ( $p=0.028$  and  $p=0.016$ ).<sup>[1]</sup> As the degree of spermatozoa with damaged DNA increases (>30-40%), the likelihood of natural pregnancy drops.<sup>[2,3]</sup> Sperm DNA of poor quality has been shown to impair fertilization by *in vitro* studies. In fact, when the degree of sperm with damaged DNA is above 4% in patients receiving *in vitro* fertilization (IVF), the fertilization rates drop from 58% to 38% ( $p<0.05$ ).<sup>[4]</sup> The importance of these data during intracytoplasmic sperm injection (ICSI) emerges from the fact that damaged DNA may not inhibit fertilization, thereby allowing the formation of embryos with this defective genetic material.<sup>[5,6]</sup> Therefore, it is important to know the degree of spermatozoa with damaged DNA in the ejaculate in order to predict the fertilization rate and determine the risks that may affect the embryo.

Although high degrees of sperm DNA damage are correlated with impaired seminal parameters such as reduced sperm count and motility, men with normal seminal parameters can also exhibit high degrees of sperm DNA damage.

In this study, we compared the results of conventional semen analyses and sperm DNA damage assay in 399 patients diagnosed with male factor infertility in the infertility clinic.

## Material and methods

This study was performed with patients who applied to Ankara University Infertility Research Center between 2010 and 2014. Men who failed to have a child despite 1 year of attempt were included in the study. Azoospermic men were excluded from the study. The sperm analyses were performed based on the WHO 2010 criteria. Seminal fluid samples were collected following a fasting period of 72 hours; a sperm count  $\geq 15$  million, forward motility  $\geq 32\%$ , and Diff-Quik morphology  $\geq 4\%$  were recognized as normal.

Ankara University Faculty of Medicine's Institutional Ethical Committee approved the study and the written informed consent was obtained from the entire subject who participated in the study.

## Analysis of total sperm DNA damage

Total Sperm DNA Damage Count (TDC), developed by Singh et al.<sup>[7]</sup>, was analyzed by alkaline assay with minor modifications. First, sperm cells were washed, followed by the mixing of 10  $\mu$ L of fresh sperm cell suspension (around 20.000 cells) with 80  $\mu$ L of 0.7% low-melting-point agarose (LMA) (Sigma) in PBS at 37°C. Subsequently, 80  $\mu$ L of this mixture was layered onto slides that had been previously coated with 1.0% hot (60°C) normal melting point agarose (NMA), covered with a coverslip at 4°C for at least 5 min to allow the agarose to solidify. After removing the coverslips, the slides were embedded in freshly prepared cold (4°C) lysing solution (2.5 M NaCl, 100 mM EDTA-<sup>2</sup>Na; 10 mM Tris-HCl, pH 10-10.5; 1% Triton X-100 and 10% DMSO added just before analysis) and 15  $\mu$ L proteinase K for at least 1 h. Slides were then immersed in freshly prepared alkaline electrophoresis buffer (0.3 mmol/L NaOH and 1 mmol/L Na<sub>2</sub>EDTA, pH >13) at 4°C for unwinding (40 min), and then subjected to electrophoresis (25 V/300 mA, 25 min). All of the above steps were conducted without applying direct light in order to prevent additional DNA damage. After electrophoresis, the slides were stained with ethidium bromide, covered with a coverslip, and analyzed using a fluorescence microscope (Olympus BX51). The images of 100 randomly chosen nuclei were analyzed visually from each subject, as described elsewhere. Each image was classified according to the intensity of the fluorescence in the comet tail and was given a value of 0, 1, 2, 3, or 4 (from undamaged class 0 to maximally damaged class 4), so that the total scores ranged between 0 and 400 arbitrary units (AU) (Figure 1).<sup>[8]</sup>

## Statistical analysis

The study data analysis was performed using Statistical Package for the Social Sciences for Windows, version 15.0 (SPSS, Inc., Chicago, IL, USA). The statistical analyses between compared groups were performed by two-sample t-test. Each p value <0.05 was considered statistically significant. Pearson and Spearman correlation coefficients were calculated between seminal parameters and TDC analysis. Reliability of the scoring systems was tested with the correlation coefficient ( $r$ ).

## Results

A total of 399 consecutive patients were included in the study. The mean age was 28 years (range: 23-42 years). Seminal parameters and TDC results were compared in these patients. Of the 399 infertile patients, mean sperm count was 45.7  $10^6$ /mL and forward motility was 37%. Of the 392 patients, mean Kruger morphology was 2.37% (Table 1).

In our study, seminal parameters were categorized based on the WHO 2010 criteria. Sperm count was classified as <15 million and  $\geq 15$  million groups. When sperm count and TDC were com-




Table 1. Mean sperm values		
Seminal parameters	n	Mean (min-max)
Sperm count x 10 <sup>6</sup> /mL	399	45.70 (0-280)
Forward motility (%)	377	37.00 (0-92)
Morphology (4%)	392	2.37 (0-21)

pared, 249 patients with normozoospermia, a sperm count  $\geq 15$  million, had a TDC of 191 AU, while 144 patients with oligozoospermia,  $< 15$  million, had a TDC value of 226 AU ( $p < 0.001$ ) (Figure 2a).

The patients were categorized with regard to sperm morphology, as well:  $< 4\%$  and  $\geq 4\%$  groups. A total of 294 patients with a morphology rate  $< 4\%$  had a TDC of 210 AU, while 97 patients with normal morphology ( $\geq 4\%$ ) exhibited a TDC of 186 AU ( $p < 0.001$ ) (Table 2).

In the present study, we observed the relation of TDC with sperm count, motility, and morphology. According to the correlation analysis, all three seminal parameters showed a negative correlation with TDC and the correlation coefficients were -0.301, -0.248 and -0.207, respectively (Table 2) (Figure 2b).

Forward motility rate was categorized in two groups:  $> 32\%$  and  $\leq 32$ . A total of 145 patients with a motility rate  $> 32\%$  had a TDC of 216 AU, whereas the patient group with a motility rate  $\leq 32$  showed a TDC of 194 AU ( $p < 0.001$ ) (Figure 2c). Comparison of the TDC values was also done between normozoospermic infertile men and infertile men with Oligoasthenoteratozoospermia (OAT). TDC was found significant-

ly higher in OAT group than those of the normozoospermic group; as 211 and 176 AU, respectively (Table 3) (Figure 2d).

## Discussion

Current evidences reveal that the most influential cause of sperm DNA damage is oxidative stress. Such events are accompanied by impaired spermiogenesis and increased ROS production due to spermatozoa. Impaired spermiogenesis is probably caused by oxidative stress leading to disrupted protein translation. Thus, spermatozoa differentiate into germ cells with certain deficiencies. Insufficient protamination also causes poor compaction of DNA which renders it vulnerable to oxidative stress. Finally, defective sperm cells may prematurely enter apoptosis, which is accompanied by the activation of mitochondrial ROS generation. As a consequence of this cascade, in addition to the disruption of sperm production, the spermatozoa lose their motility and exhibit oxidative DNA damage that leads to the creation of DNA fragmentation.<sup>[9]</sup>

Our results indicate that DNA damage is significantly higher in infertile men with OAT when compared with those of the normozoospermic men. Many studies have been performed to investigate the relation between sperm DNA damage and conventional seminal parameters. Most of these studies have yielded results indicating an inverse correlation of DNA damage with seminal parameters including sperm count, motility, and morphology.<sup>[10-12]</sup> In our study, among patients with a mature sperm count  $\geq 15 \times 10^6/\text{mL}$ , the TDC was 191 AU, whereas those with a mature sperm count  $< 15 \times 10^6/\text{mL}$  had a TDC of 226 AU (Figure 2a). The low sperm count was associated with impaired spermatogenesis; the oxidative stress generated by increasing the




Figure 2. a-d. The graphics of comparison between total sperm DNA damage count and Kruger morphology (a) sperm count (b) forward motility (c) in infertile men. Comparison of the total sperm DNA damage values in NS infertile men (All seminal parameters are in normal ranges) and infertile men with OAT (d)

ROS level hinders the maturation of germ cells by inducing DNA damage in the germ cells and accelerating apoptosis via cytochrome C and caspase 9 and caspase 3.<sup>[13]</sup>

Abortive apoptotic pathway has been proposed as an important etiology for the DNA breaks in the spermatozoa of infertile patients.<sup>[14]</sup> In the testes, normally, apoptosis prevents the overproduction of germ cells and selectively destroys injured germ cells. However, Sertoli cells are only able to support a limited number of germ cells in the testis.<sup>[15]</sup> There is an optimal germ cell to Sertoli cell ratio for the spermatogenesis to proceed normally. During this developmental process, excess immature germ cells are removed by apoptosis. If this process is impaired, the germ cell to Sertoli cell ratio is perturbed which can in turn interrupt the normal spermatogenic cascade by the result of an abortive apoptotic process. Indeed, men exhibiting deficiencies

in their semen profile often possess a large number of Fas bearing spermatozoa which leads to cell death through apoptosis.<sup>[16]</sup> This fact prompts the suggestion that these dysfunctional cells are the product of an incomplete apoptotic cascade. In case of exposure of human spermatozoa to hydrogen peroxide, apoptotic cascade has been shown to be triggered by the activation of caspase 3,<sup>[17]</sup> which may explain the association of high DNA damage and low mature spermatozoa count in our series. It has also been demonstrated that the activation of apoptotic cascade results in the stimulation of mitochondrial free radical generation.<sup>[14]</sup> Mitochondrial source of ROS can make an additional contribution to the induction of oxidative stress and DNA damage in spermatozoa.

In our study, sperm morphology and TDC values showed a significant correlation; patients with a morphology rate  $< 4\%$  had

**Table 2. Correlation between total sperm DNA damage count and seminal parameters in infertile men**

Seminal parameters		n (%)	Total sperm DNA damage count (0-400 AU)	Coefficient	p
Count 10 <sup>6</sup> /mL	<15	144 (36.64)	226±54.39	r= -0.301	<0.001
	≥15	249 (63.36)	191±48.20		
Forward motility, %	<32	145 (39.18)	216±55.13	r= -0.248	<0.001
	≥32	225 (60.82)	194±49.38		
Morphology, %	<4	294 (75.20)	210±50.12	r= -0.207	<0.001
	≥4	97 (24.8)	186±56.46		

**Table 3. Comparison of the total sperm DNA damage values in normozoospermic infertile men and infertile men with OAT**

Seminal parameters	n (%)	Total sperm DNA damage count (0-400 AU)	p
Normozoospermia*	67 (83)	176.7±44.4	<0.001
OAT**	332 (17)	211.8±53.4	<0.001

OAT: oligoasthenoteratozoospermia  
 \*All seminal parameters are in normal ranges  
 \*\*Oligoasthenoteratozoospermia; count <15x10<sup>6</sup>/mL, forward motility <32%, morphology <4%

a TDC of 186 AU, while patients exhibiting a morphology rate ≥4% showed a TDC of 210 AU (Figure 2b). In the literature, a significant association has been reported between normal sperm morphology rate and DNA fragmentation.<sup>[18]</sup>

Particularly, the incidence of serious morphologic disorders such as abnormally small heads, megaloccephaly, severe tail abnormalities, or globozoospermia has a significant correlation with abnormal chromatin structure and DNA strand breaks.<sup>[19]</sup> In such cases, there are also parallel increases in aneuploidy rates and unexplained recurrent pregnancy losses.<sup>[20]</sup>

The shaping of the sperm head is probably related to the degree of chromatin condensation as well as developmental factors.<sup>[21]</sup> It has been suggested that the shaping of sperm heads may depend on the specific patterns of assembly of DNA-protein complexes established during chromatin condensation in the nuclei.<sup>[22]</sup> On the other hand, nuclear shape has been linked to vacuoles and heterogeneity in chromatin condensation.<sup>[23]</sup> During spermiogenesis phase, last stage of spermatogenesis, significant rearrangements take place in the sperm nucleus during the transition from round spermatids to spermatozoa<sup>[24]</sup> condensation of nuclear chromatin, formation of the flagellum apparatus, and development of the acrosomal cap. The most prevalent rearrangement is the reshaping of the nucleus. Within the nucleus, the chromatin granules undergo progressive condensation due to

replacement of the transitional proteins by protamins. Protamins form a fine homogeneous material that uniformly fills the entire sperm nucleus. In light of all those events, the positive correlation between sperm DNA damage and impaired sperm morphology may be associated with the fact that sperm DNA damage leads to impairment in the sperm chromatin structure.

In our study, the degree of DNA damage and sperm motility impairment showed a positive correlation. In cases with a motility rate <32%, TDC was 216 AU, whereas patients with a motility rate ≥32% showed a TDC of 194 AU (Figure 2c). It is a well known fact that oxidative stress, which causes sperm DNA damage, induces peroxidation in the lipid-containing sperm plasma membrane via ROS, thereby leading to structural and functional damage in the spermatozoa.<sup>[25]</sup> At the same time, ROS can impair the sperm motility by damaging the axonemal structure or reducing the intracellular adenosine triphosphate (ATP).<sup>[26]</sup> Therefore, increased oxidative stress is a very effective factor in the impairment of sperm motility and the development of asthenozoospermia. The most cytotoxic oxygen metabolite among ROS products has been shown to be hydrogen peroxide. Indeed, when Percoll-separated spermatozoa are treated with hydrogen peroxide, there is a progressive decrease, leading to a complete arrest in the sperm flagellar beat frequency. However, once demembranated, ROS-immobilized spermatozoa regain motility.<sup>[27]</sup> ROS-related increases in the motility can be prevented with catalase.<sup>[28]</sup> The impairment in sperm motility has also been explained by apoptosis. It has been suggested that oxidative stress causes the generation of spermatozoa with poorly remodelled chromatin. These defective cells have a tendency to enter in an apoptotic pathway associated with motility loss.<sup>[14]</sup>

In experimental studies, a dramatic increase in DNA fragmentation has been observed after exposure to ROS products such as xanthine and xanthine oxidase. The results demonstrate that oxidative stress impairs sperm motility, as well as DNA integrity, by altering intracellular signalling pathways through changes in tyrosine phosphorylation and acid phosphatase activity.<sup>[29]</sup> In other studies, oxidative stress has been shown to promote a dose-dependent increase in tyrosine nitration and S-gluthio-



nylation, while altering motility and the ability of spermatozoa to undergo capacitation.<sup>[30]</sup>

Another cause of the impairment in sperm motility is the disruption in apoptotic balance due to oxidative stress. Apoptotic mediators such as caspases have been shown to occur at high levels in sperms with impaired motility.<sup>[31,32]</sup> A significant positive correlation between caspase activity in the sperm midpiece and the DNA fragmentation has been shown in the low motility fractions of patients, suggesting that caspase-dependent apoptotic mechanisms can originate in the cytoplasmic droplet or within the mitochondria, and function in the nucleus.<sup>[31]</sup>

The most frequently used tests to assess sperm DNA damage are the Sperm Chromatin Structure Assay (SCSA), the single-cell gel electrophoresis assay (COMET), the terminal deoxynucleotidyl transferase dUTP nick-end labeling (TUNEL) assay and the sperm chromatin dispersion (SCD) test. Because of its high precision and sensitivity in the determination of cell death and DNA damage, we prefer comet assay. The comet assay is a single cell gel electrophoresis of immobilized sperm which involves their encapsulation in agarose, followed by lysis and electrophoresis.<sup>[7]</sup> Although the technique is often tedious and may damage the cell membrane changing the distribution of the cell population of live, apoptotic or necrotic cells, it can provide more specific information about the extent and heterogeneity of DNA damage compared to TUNEL staining.<sup>[33]</sup> The TUNEL assay is also impeded by the highly compacted nature of sperm chromatin and necessitate the introduction of a chromatin decondensation step, prior to commencement of the assay in order to avoid potential artefacts.<sup>[34]</sup> On the other hand, both the TUNEL assay and the SCD test have been reported to require higher levels of DNA and chromatin damage before they can detect structural changes, indicating that they have lower sensitivity than the SCSA and the comet assay.<sup>[35]</sup>

In conclusion, as far as the comet assay can detect, high degrees of sperm DNA damage are accompanied by significant impairment in all seminal parameters. However, further studies including larger samples, homogenous patient groups, and comparison of different methods are required to reach a definitive conclusion.

**Ethics Committee Approval:** Ethics committee approval was obtained (approval number: 154-4951).

**Informed Consent:** Written informed consent was obtained from patients who participated in this study.

**Peer-review:** Externally peer-reviewed.

**Author Contributions:** Concept - O.S.A., K.A.; Design - O.S.A., K.A.; Supervision - O.S.A., A.S.; Funding - O.S.A., A.S., K.A.; Materials - O.S.A., A.S.; Data Collection and/or Processing - K.A.,

O.S.A., Y.Y., F.K.; Analysis and/or Interpretation - O.S.A., Y.Y., F.K., K.A.; Literature Review - O.S.A., Y.Y., F.K.; Writer - O.S.A.; Critical Review - K.A., A.S.

**Conflict of Interest:** The authors declared no conflict of interest.

**Financial Disclosure:** This research was supported by Ankara University Scientific Research Division (project Number: 10B3330028). The research facilities for this project were provided by Ankara University School of Medicine, Department of Medical Biology.

**Etik Komite Onayı:** Bu çalışma için etik komite onayı alınmıştır (onay numarası: 154-4951).

**Hasta Onamı:** Yazılı hasta onamı bu çalışmaya katılan hastalardan alınmıştır.

**Hakem Değerlendirmesi:** Dış bağımsız.

**Yazar Katkıları:** Fikir - O.S.A., K.A.; Tasarım - O.S.A., K.A.; Denetleme - O.S.A., A.S.; Kaynaklar - O.S.A., A.S., K.A.; Malzemeler - O.S.A., A.S.; Veri toplanması ve/veya işlemesi K.A., O.S.A., Y.Y., F.K.; Analiz ve/veya yorum - O.S.A., Y.Y., F.K., K.A.; Literatür taraması - O.S.A., Y.Y., F.K.; Yazıyı yazan - O.S.A.; Eleştirel inceleme - K.A., A.S.

**Çıkar Çatışması:** Yazarlar çıkar çatışması bildirmemişlerdir.

**Finansal Destek:** Tıbbi Biyoloji bölümünün bu projesi Ankara Üniversitesi Bilimsel Araştırmalar tarafınca için desteklenmiştir (proje numarası: 10B3330028).

## References

1. Zini A, Bielecki R, Phang D, Zenzes MT. Correlations between two markers of sperm DNA integrity, DNA denaturation and DNA fragmentation, in fertile and infertile men. *Fertil Steril* 2001;75:674-7. [\[CrossRef\]](#)
2. Evenson DP, Jost LK, Marshall D, Zinaman MJ, Clegg E, Purvis K, et al. Utility of the sperm chromatin structure assay as a diagnostic and prognostic tool in the human fertility clinic. *Hum Reprod* 1999;14:1039-49. [\[CrossRef\]](#)
3. Spano M, Bonde JP, Hgjollund HI, Kolstad HA, Cordelli E, Leter G. Sperm chromatin damage impairs human fertility. The Danish first pregnancy planner study team. *Fertil Steril* 2000;73:43-50.
4. Host E, Lindenberg S, Smidt-Jensen S. The role of DNA strand breaks in human spermatozoa used for IVF and ICSI. *Acta Obstet Gynecol Scand* 2001;79:559-63. [\[CrossRef\]](#)
5. Lopes SM, Sun JG, Jurisicova A, Meriano JB, Casper RF. Sperm deoxyribonucleic acid fragmentation is increased in poor-quality semen samples and correlates with failed fertilization in intracytoplasmic sperm injection. *Fertil Steril* 1998;69:528-32. [\[CrossRef\]](#)
6. Ahmadi A, Ng SC. Fertilizing ability of DNA-damaged spermatozoa. *J Exp Zool* 1999;284:696-704. [\[CrossRef\]](#)
7. Singh NP. Microgel electrophoresis of DNA. In: *Technologies for Detection of DNA Damage and Mutations*. New York: Plenum Press; 1996. p.3-24. [\[CrossRef\]](#)
8. Jalszynski P, Kujawski M, Czub-Swierczek M, Markowska J, Szyfter K. Bleomycin- induced DNA damage and its removal in

- lymphocytes of breast cancer patients studied by comet assay. *Mutat Res* 1997;385:223-33. [\[CrossRef\]](#)
9. Aitken RJ, Curry BJ. Redox regulation of human sperm function: from the physiological control of sperm capacitation to the etiology of infertility and DNA damage in the germ line. *Antioxid Redox Signal* 2011;14:367-81. [\[CrossRef\]](#)
  10. Acharyya S, Kanjilal S, Bhattacharyya AK. Does human sperm nuclear DNA integrity affect embryo quality? *Indian J Exp Biol* 2005;43:1016-22.
  11. Wyrobek AJ, Eskenazi B, Young S, Arnheim N, Tiemann-Boege I, Jabs EW, et al. Advancing age has differential effects on DNA damage, chromatin integrity, gene mutations, and aneuploidies in sperm. *P Natl Acad Sci USA* 2006;103:9601-6. [\[CrossRef\]](#)
  12. Evgeni E, Lymberopoulos G, Gazouli M, Asimakopoulos B. Conventional semen parameters and DNA fragmentation in relation to fertility status in a Greek population. *Eur J Obstet Gynecol Reprod Biol* 2015;188:17-23. [\[CrossRef\]](#)
  13. Said TM, Paasch U, Glander HJ, Agarwal A. Role of caspases in male infertility. *Hum Reprod Update* 2004;10:39-51. [\[CrossRef\]](#)
  14. Aitken RJ, Koppers AJ. Apoptosis and DNA damage in human spermatozoa. *Asian J Androl* 2011;13:36-42. [\[CrossRef\]](#)
  15. Sinha Hikim AP, Swerdloff RS. Hormonal and genetic control of germ cell apoptosis in the testis. *Rev Reprod* 1999;4:38-47. [\[CrossRef\]](#)
  16. Sakkas D, Mariethoz E, Manicardi G, Bizzaro D, Bianchi PG, Bianchi U. Origin of DNA damage in ejaculated human spermatozoa. *Rev Reprod* 1999;4:31-7. [\[CrossRef\]](#)
  17. Lozano GM, Bejarano I, Espino J, González D, Ortiz A, García JF, et al. Relationship between caspase activity and apoptotic markers in human sperm in response to hydrogen peroxide and progesterone. *J Reprod Dev* 2009;55:615-21. [\[CrossRef\]](#)
  18. Menkveld R, Holleboom CA, Rhemrev JP. Measurement and significance of sperm morphology. *Asian J Androl* 2011;13:59-68. [\[CrossRef\]](#)
  19. Ombelet W, Wouters E, Boels L, Cox A, Janssen M, Spiessens C, et al. Sperm morphology assessment: diagnostic potential and comparative analysis of strict or WHO criteria in a fertile and a subfertile population. *Int J Androl* 1997;20:367-72. [\[CrossRef\]](#)
  20. Carrell DT, Wilcox AL, Lowy L, Peterson CM, Jones KP, Erickson L, et al. Elevated sperm chromosome aneuploidy and apoptosis in patients with unexplained recurrent pregnancy loss. *Obstet Gynecol* 2003;101:1229-35. [\[CrossRef\]](#)
  21. Pedersen H, Fawcett DW. Biology of spermatozoa In *Human semen and fertility regulation in men*. Saint Louis:The CV Mosby Co; 1976. p.65.
  22. Fawcett DW, Anderson WA, Phillips DM. Morphogenetic factors influencing the shape of the sperm head. *Dev Biol* 1971;26:220-51. [\[CrossRef\]](#)
  23. Curry MR, Watson PF. Sperm structure and function. In: *Gametes the Spermatozoon*. Cambridge:Cambridge University Press; 1995. p.45.
  24. Hafez ESE, Hafez SD. Atlas of clinical andrology. Oxon, UK:Taylor and Francis Group; 2005. p.15. [\[CrossRef\]](#)
  25. Agarwal A, Said TM. Oxidative stress, DNA damage and apoptosis in male infertility: a clinical approach. *BJU Int* 2005;95:503-7. [\[CrossRef\]](#)
  26. Tsunoda S, Kawano N, Miyado K, Kimura N, Fujii J. Impaired fertilizing ability of superoxide dismutase 1-deficient mouse sperm during in vitro fertilization. *Biol Reprod* 2012;87:121. [\[CrossRef\]](#)
  27. de Lamirande E, Gagnon C. Reactive oxygen species and human spermatozoa. I. Effects on the motility of intact spermatozoa and on sperm axonemes. *J Androl* 1992;13:368-78.
  28. Martínez-Pastor F, Aisen E, Fernández-Santos MR, Estes MC, Maroto-Morales A, García-Alvarez O, et al. Reactive oxygen species generators affect quality parameters and apoptosis markers differently in red deer spermatozoa. *Reproduction* 2009;137:225-35. [\[CrossRef\]](#)
  29. Gazo I, Shaliutina-Kolešová A, Dietrich MA, Linhartová P, Shaliutina O, Cosson J. The effect of reactive oxygen species on motility parameters, DNA integrity, tyrosine phosphorylation and phosphatase activity of common carp (*Cyprinus carpio* L.) spermatozoa. *Mol Reprod Dev* 2015;82:48-57. [\[CrossRef\]](#)
  30. Morielli T, O'Flaherty C. Oxidative stress impairs function and increases redox protein modifications in human spermatozoa. *Reproduction* 2015;149:113-23. [\[CrossRef\]](#)
  31. Weng SL, Taylor SL, Morshedi M, Schuffner A, Duran EH, Beebe S, et al. Caspase activity and apoptotic markers in ejaculated human sperm. *Mol Hum Reprod* 2002;8:984-91. [\[CrossRef\]](#)
  32. Taylor SL, Weng SL, Fox P, Duran EH, Morshedi MS, Oehninger S, et al. Somatic cell apoptosis markers and pathways in human ejaculated sperm: potential utility as indicators of sperm quality. *Mol Hum Reprod* 2004;10:825-34. [\[CrossRef\]](#)
  33. Archana M, Bastian Yogesh TL, Kumaraswamy KL. Various methods available for detection of apoptotic cells a review. *Indian J Cancer* 2013;50:274-83. [\[CrossRef\]](#)
  34. Aitken RJ, Bronson R, Smith TB, De Iulius GN. The source and significance of DNA damage in human spermatozoa; a commentary on diagnostic strategies and straw man fallacies. *Mol Hum Reprod* 2013;19:475-85. [\[CrossRef\]](#)
  35. Pérez-Cereales S, Miranda A, Gutiérrez-Adán A. Comparison of four methods to evaluate sperm DNA integrity between mouse caput and cauda epididymidis. *Asian J Androl* 2012;14:335-7. [\[CrossRef\]](#)